arXiv:cond-mat/9906078v1 [cond-mat.dis-nn] 5 Jun 1999

International Journal of Modern Physics C, Vol. 0, No. 0 (1999) 000–000
© World Scientific Publishing Company

AN INVESTIGATION OF SITE-BOND PERCOLATION ON MANY LATTICES

YURIY YU. TARASEVICH
Faculty of Physics and Mathematics, Astrakhan State Pedagogical University
Astrakhan 414056, Russia
E-mail: vt@astranet.ru

and

STEVEN C. VAN DER MARCK*
SIEP Research and Technical Services, P.O. Box 60
2280 AB Rijswijk, The Netherlands
E-mail: steven.vandermarck@wxs.nl

Received (received date)
Revised (revised date)

A calculation of site-bond percolation thresholds in many lattices in two to five dimensions is presented. The line of threshold values has been parametrized in the literature, but we show here that there are strong deviations from the known approximate equations. We propose an alternative parametrization that lies much closer to the numerical values.

Keywords: Mixed Percolation; Critical Curve; Lattice

1. Introduction

Site-bond percolation is a natural generalization of pure site percolation and pure bond percolation. The generalization allows both sites and bonds to be randomly occupied, with probabilities p_s and p_b respectively, in the case of random percolation. In particular, if $p_b = p_s$, site-bond percolation reduces to a pure site or pure bond percolation on a decorated lattice. It is also possible to introduce a correlation between the occupation probabilities of adjacent sites and bonds, which is usually referred to as correlated percolation. Site-bond percolation has many applications in different fields. For example, it was used to describe the sol-to-gel transition (gelation) of polymers,[1] the spreading of an infection,[2] etc.

In site-bond percolation one looks for clusters of occupied sites, connected to each other by occupied bonds. When there is a cluster that connects one side of the system with the other, the system is said to percolate. First mentioned by Frisch and Hammersley,[3] the model was investigated at first by Hoshen by means of Monte Carlo simulations.[4] Agrawal et al.[5] showed, using a series method, that the critical

*Present address: EDS, P.O. Box 406, 2260 AK Leidschendam, The Netherlands

exponents of pure site percolation are also valid for site-bond percolation. Nakanishi and Reynolds[6] confirmed this conclusion through a position-space renormalization group study. They also found that the boundary between percolation and non-percolation in the (p_b, p_s) plane is governed by a single fixed point. Hammersley[2] proved for a partially directed graph a theorem. The theorem yields the inequality

$$P(p_s p_b, 1) \leq P(p_s, p_b) \leq P(1, p_s p_b), \tag{1}$$

where $P(p_s, p_b)$ is the percolation probability (i.e. the probability that a single source site is connected to an infinite set of other sites). The inequality Eq. (1) gives reasonably sharp bounds for the mixed percolation probability. For a Bethe lattice ($p_{cb} = p_{cs} = p_c$, where p_{cs} is the threshold for pure site percolation, and p_{cb} is the one for pure bond percolation) we can get

$$p_s p_b = p_c \tag{2}$$

The shape of the boundary between percolation and non-percolation was subsequently studied by Yanuka and Englman.[7] They proposed the following equation for the 'critical curve' in the (p_b, p_s) plane.

$$\frac{\log p_s}{\log p_{cs}} + \frac{\log p_b}{\log p_{cb}} = 1, \tag{3}$$

The equation satisfies the inequality Eq. (1) and gives the correct limit Eq. (2) for a Bethe lattice. The equation defines a line in the (p_b, p_s) plane, starting at $(p_{cb}, 1)$ and ending at $(1, p_{cs})$, as is shown in Figure 1. Yanuka and Englman showed plots of Eq. (3) together with simulation results for (p_b, p_s) for the square, triangular, simple cubic and face centered cubic (fcc) lattices. All their simulation results lie remarkably close to the line defined by Eq. (3). It is not clear from their paper whether the data of e.g. the fcc lattice, which lie visibly below the line given by Eq. (3), are accurate enough to conclude that there is a discrepancy.

Tretyakov and Inui[8] studied the same critical curve, but now for directed percolation. Their simulation results were accurate enough to conclude that deviations in the range $10^{-3} - 10^{-4}$ occur. This raises the question whether such deviations also occur for 'regular' site-bond percolation.

Therefore we have undertaken a systematic study of the threshold values (p_b, p_s) for random site-bond percolation in many lattices. The results, which are presented in Section , show many deviations from the Yanuka-Englman line of Eq. (3). The deviations are largest for lattices with a low ratio of $\alpha = p_{cb}/p_{cs}$, i.e. lattices for which the bond percolation thresholds is much lower than the site percolation threshold. In Section 2 we discuss other approximative equations for the critical line (p_b, p_s).

2. Numerical results

All numerical results were obtained with the programs described in Ref.[9a]. For

[a]The programs can be obtained from one of the authors by e-mail at steven.vandermarck@wxs.nl



Fig. 1. The percolation threshold pairs (p_s, p_b) for many different lattices in various dimensions. The points are from the simulations (see Tables 1–4), the lines are from Eq. (3).

Table 1. The results for the site-bond percolation thresholds for several lattices in two dimensions. Error estimates concerning the last digit(s) are indicated between brackets.

	square		triangular		Kagomé	
p	$p_b(p_s=p)$	$p_s(p_b=p)$	$p_b(p_s=p)$	$p_s(p_b=p)$	$p_b(p_s=p)$	$p_s(p_b=p)$
0.95	0.5334(3)	0.6149(3)	0.3732(3)	0.5152(4)	0.5615(4)	0.6711(4)
0.90	0.5716(3)	0.6396(4)	0.4023(3)	0.5331(4)	0.6046(5)	0.6914(5)
0.85	0.6154(3)	0.6673(3)	0.4368(3)	0.5518(4)	0.6556(6)	0.7162(5)
0.80	0.6663(3)	0.6980(3)	0.4764(4)	0.5736(4)	0.7152(7)	0.7428(5)
0.75	0.7257(4)	0.7322(4)	0.5251(4)	0.5969(4)	0.7894(9)	0.7757(8)
0.70	0.7961(4)	0.7709(3)	0.5830(6)	0.6231(3)		
0.65	0.8807(4)	0.8157(3)	0.6561(6)	0.6541(4)		
0.60	0.9840(4)	0.8671(3)				
	octagonal		honeycomb		star-dual	
p	$p_b(p_s=p)$	$p_s(p_b=p)$	$p_b(p_s=p)$	$p_s(p_b=p)$	$p_b(p_s=p)$	$p_s(p_b=p)$
0.95	0.3472(4)	0.5133(7)	0.6926(5)	0.7275(5)	0.2796(4)	0.5036(5)
0.90	0.3744(5)	0.5276(9)	0.7377(5)	0.7610(5)	0.3022(5)	0.5101(5)
0.85	0.4087(5)	0.5442(7)	0.7890(5)	0.7986(5)	0.3296(4)	0.5188(5)
0.80	0.4449(6)	0.5620(7)	0.8481(5)	0.8401(5)	0.3615(5)	0.5285(5)
0.75	0.4912(7)	0.5816(7)			0.4006(4)	0.5417(5)
0.70	0.5481(10)	0.6052(7)			0.4499(5)	0.5572(5)
0.65	0.6213(24)				0.5133(4)	0.5763(5)
0.60					0.5968(5)	0.5984(5)
	puzzle		direct		low-pb	
p	$p_b(p_s=p)$	$p_s(p_b=p)$	$p_b(p_s=p)$	$p_s(p_b=p)$	$p_b(p_s=p)$	$p_s(p_b=p)$
0.95	0.5693(5)	0.4495(4)	0.5677(4)	0.4433(4)	0.2166(5)	0.5003(5)
0.90	0.5901(5)	0.4849(5)	0.5877(4)	0.4780(4)		
0.85	0.6124(4)	0.5247(5)	0.6091(4)	0.5183(5)	0.2554(5)	0.5048(5)
0.80	0.6371(5)	0.5720(5)	0.6343(4)	0.5648(4)		
0.75	0.6655(5)	0.6283(5)	0.6602(4)	0.6199(5)	0.3110(5)	0.5156(5)
0.70	0.6976(4)	0.6952(5)	0.6913(4)	0.6876(5)		
0.65	0.7328(5)	0.7766(5)	0.7263(4)	0.7692(5)	0.4020(5)	0.5334(5)
0.60			0.7674(5)	0.8712(7)		
0.55					0.5644(5)	
0.45					0.6149(5)	
0.35					0.6997(6)	

Table 2. The results for the site-bond percolation thresholds for several lattices in three dimensions. Error estimates concerning the last digit(s) are indicated between brackets.

	sc		fcc		hcp	
p	$p_b(p_s=p)$	$p_s(p_b=p)$	$p_b(p_s=p)$	$p_s(p_b=p)$	$p_b(p_s=p)$	$p_s(p_b=p)$
0.95	0.2637(3)	0.3244(3)	0.1273(3)	0.2049(5)	0.1272(3)	0.2052(3)
0.90	0.2806(3)	0.3385(3)	0.1355(3)	0.2124(4)	0.1361(3)	0.2125(3)
0.85	0.2991(3)	0.3539(3)	0.1453(3)	0.2202(4)	0.1456(3)	0.2214(3)
0.80	0.3208(3)	0.3716(3)	0.1560(3)	0.2291(5)	0.1559(3)	0.2306(3)
0.75	0.3452(3)	0.3912(3)	0.1679(3)	0.2391(5)	0.1688(3)	0.2404(3)
0.70	0.3735(3)	0.4139(3)	0.1823(3)	0.2515(5)	0.1832(3)	0.2519(3)
0.65	0.4072(3)	0.4395(3)	0.1995(4)	0.2644(5)	0.2001(3)	0.2653(3)
0.60	0.4475(3)	0.4694(3)	0.2201(4)	0.2796(6)	0.2209(3)	0.2806(3)
0.55			0.2443(5)	0.2974(6)	0.2455(3)	0.2985(3)
0.50			0.2769(5)	0.3190(6)	0.2764(3)	0.3195(3)
0.45			0.3161(6)	0.3451(6)	0.3164(3)	0.3454(3)
0.40			0.3676(8)	0.3759(6)	0.3698(3)	0.3764(3)
0.35			0.4404(11)	0.4171(7)		

	bcc		diamond	
p	$p_b(p_s=p)$	$p_s(p_b=p)$	$p_b(p_s=p)$	$p_s(p_b=p)$
0.95	0.1909(3)	0.2545(5)	0.4118(4)	0.4489(5)
0.90	0.2030(3)	0.2656(5)	0.4366(4)	0.4723(5)
0.85	0.2164(4)	0.2777(5)	0.4656(4)	0.4959(5)
0.80	0.2324(4)	0.2900(5)	0.4973(4)	0.5229(5)
0.75	0.2498(4)	0.3043(5)	0.5345(5)	0.5544(6)
0.70	0.2711(4)	0.3210(6)	0.5758(6)	0.5885(5)
0.65	0.2953(5)	0.3408(6)	0.6253(6)	0.6283(5)
0.60	0.3249(6)	0.3633(6)	0.6845(7)	0.6755(6)
0.55	0.3603(6)	0.3896(7)	0.7555(8)	0.7308(6)
0.50	0.4036(7)	0.4199(7)		
0.45	0.4578(9)	0.4562(7)		
0.40	0.5300(10)	0.5043(7)		
0.35	0.6281(13)	0.5639(7)		

Table 3. The results for the site-bond percolation thresholds for the 4-dimensional simple cubic (sc) and fcc lattices. Error estimates concerning the last digit(s) are indicated between brackets.

	4d-sc		4d-fcc	
p	$p_b(p_s = p)$	$p_s(p_b = p)$	$p_b(p_s = p)$	$p_s(p_b = p)$
0.95	0.1701(6)	0.2065(6)	0.0520(03)	0.0853(08)
0.90			0.0554(04)	0.0892(09)
0.85	0.1900(6)	0.2254(6)	0.0584(04)	0.0920(10)
0.80			0.0635(04)	0.0979(09)
0.75	0.2183(6)	0.2509(6)	0.0668(05)	0.1028(10)
0.70			0.0722(05)	0.1052(11)
0.65	0.2547(6)	0.2837(6)	0.0789(06)	0.1139(10)
0.60			0.0868(07)	0.1167(13)
0.55	0.3064(6)	0.3275(6)	0.0957(07)	0.1276(12)
0.50			0.1067(09)	0.1372(13)
0.45	0.3834(6)	0.3897(6)	0.1171(10)	0.1473(13)
0.40			0.1337(12)	0.1627(15)
0.35	0.5083(6)	0.4880(6)	0.1579(14)	0.1769(18)
0.30			0.1877(19)	0.2034(17)
0.25	0.7512(6)	0.6618(7)	0.2339(27)	

Table 4. The results for the site-bond percolation thresholds for the 5-dimensional fcc and Kagomé lattices. Error estimates concerning the last digit(s) are indicated between brackets.

	5d-fcc		5d-Kagomé	
p	$p_b(p_s = p)$	$p_s(p_b = p)$	$p_b(p_s = p)$	$p_s(p_b = p)$
0.95	0.0274(5)	0.0434(5)	0.1374(16)	0.2173(20)
0.85	0.0319(5)	0.0474(6)	0.1555(17)	0.2284(19)
0.75	0.0365(5)	0.0522(6)	0.1851(19)	0.2439(20)
0.65	0.0424(5)	0.0598(6)	0.2111(19)	0.2633(19)
0.55	0.0500(5)	0.0661(6)	0.2538(18)	0.2957(19)
0.45	0.0627(6)	0.0787(6)	0.3245(20)	0.3494(20)
0.35	0.0816(6)	0.0963(6)	0.4506(20)	0.4243(19)
0.25	0.1179(6)	0.1283(6)		

each lattice the percolation threshold was calculated for at least three values of the linear lattice size L. The percolation threshold $p_c(\infty)$ for infinite lattices can be found by fitting these results for different lattice sizes to the scaling relation

$$|p_c(L) - p_c(\infty)| \propto L^{-1/\nu}, \tag{4}$$

where the critical exponent ν has the value 4/3 in two dimensions, and 0.88, 0.68, 0.57 in 3, 4, and 5 dimensions respectively[10]. These steps are precisely the same as for pure site or pure bond percolation. The results are shown in Tables 1–4 and in Figures 1–6. We have checked that for pure site percolation, and for pure bond percolation, our program yields results equivalent to more precise literature values (see Table 5).

We verified several scaling laws in percolation, as a check on the program. First, we calculated the correlation length ξ below the percolation threshold, as a function of $p_c - p$. The correlation length scales as

$$\xi \propto (p_c - p)^{-\nu}. \tag{5}$$

The calculation was performed by first computing the 'gyration radius' R_s for a cluster of s sites. This radius is defined by

$$R_s^2 = \frac{1}{s}\sum_{i=1}^{s} |\mathbf{r}_i - \mathbf{r}_0|^2, \qquad \text{with} \qquad \mathbf{r}_o = \frac{1}{s}\sum_{i=1}^{s} \mathbf{r}_i, \tag{6}$$

where $\mathbf{r}_i$ is the position of the i^{th} site in the cluster. The correlation length ξ can then be calculated as (see Ref.[10], p. 50)

$$\xi^2 = \frac{2\sum_s R_s^2 s^2 n_s}{\sum_s s^2 n_s}, \tag{7}$$

where n_s is the mean number per lattice site of clusters of s sites. We verified numerically that the correlation length obeys the scaling relation (5). Our result for site-bond percolation on the Kagomé and diamond lattices is shown in Figure 2. A fit of the scaling relation to the data yields $\nu = 1.31 \pm 0.01$ for the Kagomé lattice, and $\nu = 0.91 \pm 0.04$ for the diamond lattice. This is within one or two standard deviations of the established values of $\nu = 4/3$ in two dimensions and $\nu = 0.88$ in three dimensions.

Also we have checked the scaling of the mean cluster size S and the strength of the infinite cluster, both as a function of $|p - p_c|$. The average cluster size S scales with the same exponent both above and below the percolation threshold:

$$S \propto |p_c - p|^{-\gamma}, \tag{8}$$

with $\gamma = 43/18$ in two dimensions, and 1.80 in three[10]. The results for site-bond percolation on the Kagomé and diamond lattices is shown in Fig. 3. The fits to the scaling relation (8) yield values for γ that are consistent with the literature values.



Fig. 2. The correlation length ξ as a function of $p_c - p$. The runs were performed at $\{p_b = 0.65,\ p_s = 0.6284(5)\}$ for the diamond lattice and $\{p_b = 0.75,\ p_s = 0.7757(8)\}$ for the Kagomé lattice.



Fig. 3. The mean cluster size S above and below the percolation threshold, as a function of $|p_c - p|$. The runs were performed at $\{p_b = 0.65,\ p_s = 0.6284(5)\}$ for the diamond lattice (solid lines) and $\{p_b = 0.75,\ p_s = 0.7757(8)\}$ for the Kagomé lattice (dashed lines).

The strength P of the infinite cluster scales as

$$P \propto (p - p_c)^{\beta}\,, \tag{9}$$

with $\beta = 5/36$ in two dimensions, and 0.41 in three[10]. The results for site-bond percolation on the Kagomé and diamond lattices is shown in Fig. 4. The fits to the scaling relation (9) yield values for β that are consistent with the literature values.



Fig. 4. The strength P of the infinite cluster as a function of $p - p_c$. The runs were performed at $\{p_b = 0.65,\ p_s = 0.6284(5)\}$ for the diamond lattice and $\{p_b = 0.75,\ p_s = 0.7757(8)\}$ for the Kagomé lattice.

3. Approximate equations

It is clear from Figure 1 that there are systematic deviations from the Yanuka-Englman line. The accuracy of the numerical values, as indicated in Tables 1–4 is easily high enough to warrant this conclusion. The Yanuka-Englman line is situated above the numerical values for all calculated examples. One can say it is an upper estimation for the real critical curve. In general it seems that the deviations are largest for lattices with a low ratio of $\alpha = p_{cb}/p_{cs}$, i.e. lattices for which the bond percolation thresholds is much lower than the site. The most clear example is the 'low-p_b' lattice (see Fig. 5), which has $p_{cs} = 0.5$ and $p_{cb} = 0.2006(3)$ (see Table 5). This lattice has been constructed from the triangular lattice in two steps, see Fig. 5. All the lattices in this figure are fully triangulated and have therefore

$p_{cs} = \frac{1}{2}$, as shown by Sykes and Essam[13]. The bond percolation threshold, however, will decrease with every step in Fig. 5, and these steps can be repeated even further. In this way one can construct lattices with arbitrarily low ratio p_{cb}/p_{cs}.



Fig. 5. The triangular lattice (left) can be modified by adding a site in the middle of each triangle (middle). The same operation once more yields the lattice on the right. All these lattice have $p_s = \frac{1}{2}$, but the bond percolation threshold decreases from left to right.

One of the most remarkable results in our calculations is the results for two pairs of lattices with very close or equal critical probabilities. One pair is the face centered cubic (fcc) lattice and the hexagonal close packed (hcp) lattice. Another pair, in two dimensions, is the direct lattice and the puzzle lattice (we use here the nicknames as in Ref. [12]). The fcc and hcp lattices, although they are topologically different, have $p_{cb}^{fcc} = p_{cb}^{hcp}$ and $p_{cs}^{fcc} = p_{cs}^{hcp}$ within the present small error margins (see Table 5). Also the critical curves we calculated for these lattices are the same within our error margin. In two dimensions, the pure site and pure bond percolation thresholds of the direct and the puzzle lattice are almost the same. Also for this pair the critical curves lie close together.

These results suggest that once the pure site and pure bond thresholds are known, the whole critical curve is fixed. In other words, there is no need to use extra parameters (such as critical exponents, dimension, or coordination number) to describe a critical curve but p_{cb} and p_{cs}. In this context it is interesting to note that in the limiting case of Bethe lattices, Eq. (2), there are also no parameters other than p_c.

One can try to find an equation for the critical curve by making use of the work of Hammersley, who derived boundaries for mixed percolation[2]

$$p_s^{up}(p_b) = \frac{p_{cs}}{p_b},$$

and

$$p_s^{down}(p_b) = \frac{p_{cb}}{p_b}.$$

Let us look for a critical relationship as a combination of these boundaries

$$p_s(p_b) = C_1(p_b) \cdot p_s^{down}(p_b) + C_2(p_b) \cdot p_s^{up}(p_b).$$

Take the functions $C_1(p_b)$ and $C_2(p_b)$ in the simplest form

$$C_1(p_b) = \frac{1 - p_b}{1 - p_{cb}} = \begin{cases} 1 & \text{for } p_b = p_{cb} \\ 0 & \text{for } p_b = 1 \end{cases},$$

and

$$C_2(p_b) = \frac{p_b - p_{cb}}{1 - p_{cb}} = \begin{cases} 1 & \text{for } p_b = 1 \\ 0 & \text{for } p_b = p_{cb} \end{cases} .$$

After some transformation we have

$$p_b\,(p_s + A) = B, \tag{10}$$

where

$$A = \frac{p_{cb} - p_{cs}}{1 - p_{cb}}, \tag{11}$$

$$B = p_{cb}\frac{1 - p_{cs}}{1 - p_{cb}}. \tag{12}$$

The relationship Eq. (10) gives almost the same results as Eq. (3) provided p_b and p_s are close (square, simple cubic, octagonal, etc.). In these cases both curves and the numerical points are in good agreement with one another. In other cases Eq. (10) gives much better approximations than Eq. (3) (see Figure 6).



Fig. 6. The percolation threshold pairs (p_s, p_b) for many different lattices in various dimensions. The points are from the simulations (see Tables 1–4), the lines are from Eq. (10).

We examined a number of different relationships with 2 and 3 parameters to describe the numerical results. Although the results were excellent for some lattices, we could not find a relationship suitable for all lattices.

Hence the problem to find the real law for critical probabilities is still unsolved. We hope this work with the large collection of high precision numerical results and the new estimation of critical probabilities will assist to solve the problem.

Acknowledgements

One of the authors (Yu.Yu. Tarasevich) thanks Max-Plank-Institut für Physik Komplexer Systeme for the possibility to work in the library during a seminar and Prof. Amnon Aharony for discussions.

Table 5. The site and bond percolation thresholds used in the article, together with the references to the literature.

lattice	p_s [Ref.]		p_b [Ref.]	
honeycomb	0.697 043 (2)	[12]	0.652 703 ...	[13]
Kagomé	0.652 703 ...	[13]	0.524 405 3(3)	[14]
square	0.592 746 0(5)	[15]	0.5	[16]
triangular	0.5	[13,16]	0.347 296 ...	[13]
octagonal	0.5	[13]	0.3230(2)	[17,11]
star-dual	0.5	[13]	0.2593(2)	[11]
low-p_b	0.5	[13]	0.2006(3)	
puzzle	0.550806(2)	[12]	0.4142(2)	
direct	0.550213(2)	[12]	0.4195(2)	
simple cubic	0.311 608 0(3)	[18]	0.248 812 6(5)	[19]
bcc	0.245 961 5(10)	[18]	0.180 287 5(10)	[19]
fcc	0.199 236 5(10)	[18]	0.120 163 5(10)	[19]
hcp	0.199 24(5)	[17]	0.120 15(5)	[17]
diamond	0.4301(2)	[9]	0.3893(2)	[17]
4d-cubic	0.196901(5)	[20]	0.1600(2)	[21]
4d-fcc	0.0842(3)	[9]	0.049(1)	[9]
5d-fcc	0.0431(3)	[9]	0.026(2)	[9]
5d-Kagomé	0.2084(4)	[22]	0.130(2)	[9]

References

1. A. Coniglio, H.E. Stanley and W. Klein, *Phys. Rev. Lett.* **42**, 518 (1979).
2. J.M. Hammersley, *Math. Proc. Camb. Phyl. Soc.* **88**, 167 (1980).
3. H.L. Frisch and J.M. Hammersley, *J. Soc. Ind. Appl. Math.* **11**, 894 (1963).
4. J. Hoshen, Univ. of Michigan, preprint (1978).
5. P. Agrawal, S. Render, P.J. Reynolds, and H.E. Stanley, *J. Phys. A: Math. Gen.* **12**, 2073 (1979).
6. H. Nakanishi and J. Reynolds, *Phys. Lett.* **A71**, 252 (1979).
7. M. Yanuka and R. Englman, *J. Phys. A: Math. Gen.* **23**, L339 (1990).
8. A. Tretyakov and N. Inui, *J. Phys. A: Math. Gen.* **28**, 3985 (1995).
9. S.C. van der Marck, *Int. J. Mod. Phys. C* **9**, 529 (1998).

10. D. Stauffer and A. Aharony, *Introduction to Percolation Theory* (Taylor and Francis, London, 1992).
11. S.C. van der Marck, to be published.
12. P.N. Suding and R.M. Ziff, cond-mat/9811416, submitted to Phys. Rev. E.
13. M.F. Sykes and J.W. Essam, *J. Math. Phys.* **5**, 1117 (1964).
14. R.M. Ziff and P.N. Suding, *J. Phys. A: Math. Gen.* **30**, 5351 (1997).
15. R.M. Ziff and B. Sapoval, *J. Phys. A: Math. Gen.* **19**, L1169 (1986); R.M. Ziff, *Phys. Rev. Lett.* **69**, 2670 (1992).
16. *H. Kesten, Comm. Math. Phys.* **74**, 41 (1980).
17. S.C. van der Marck, *Phys. Rev.* **E55**, 1514 (1997) p.1514; Erratum **56**, 3732 (1997).
18. C.D. Lorenz and R.M. Ziff, *J. Phys. A: Math. Gen.* **31**, 8147 (1998).
19. C.D. Lorenz and R.M. Ziff, *Phys. Rev.* **E57**, 230 (1998).
20. H.G. Ballesteros, L.A. Fernández, V. Martín-Mayor, A. Muñoz Sudupe, G. Parisi, and J.J. Ruiz-Lorenzo, *Phys. Lett. B* **400**, 346 (1997).
21. D.S. Gaunt and H. Ruskin, *J. Phys. A: Math. Gen.* **11**, 1369 (1978).
22. S.C. van der Marck, *J. Phys. A: Math. Gen.* **31**, 3449 (1998).